First result of the experimental search for the $2K$-capture of ^{124}Xe with the copper proportional counter[a]

Yu.M. Gavrilyuk[†], A.M. Gangapshev[†], V.V. Kazalov[†], V.V. Kuzminov[†],
S.I. Panasenko[‡], S.S. Ratkevich[‡], D.A. Tekueva[†] and S.P. Yakimenko[†]

[†]*Baksan Neutrino Observatory INR RAS, Russiaa*

[‡] *V.N.Karazin Kharkiv National University,Ukraine*

(Dated: April 23, 2014)

arXiv:1404.5530v1 [nucl-ex] 22 Apr 2014

Abstract

First result of experiment for searching of $2K$-capture of ^{124}Xe with the large-volume copper proportional counter (LPC) is given. The 12 L sample with 63.3% (44 g) of ^{124}Xe was used in measurements. The limit on the half-life of ^{124}Xe with regard to $2K(2\nu)$-capture for the ground state of ^{124}Te ($0^+ \to 0^+$, g.s.) has been found: $T_{1/2} \geq 4.67 \times 10^{20}$ y (90% C.L.). A sample with volume 52 L comprising of ^{124}Xe (10.6 L -58.6 g) and ^{126}Xe (14.1 L - 79.3 g) will used at the next step of the experiment to increase a sensitivity of 2K-caption of ^{124}Xe registration. In this case sensitivity to the investigated process will be at the level of $S = 1.46 \times 10^{21}$ y (90% C.L.) for 1 year measurement.

PACS numbers: 23.40.-s, 27.60.+j, 29.40.Cs

[a] Talk at the International Workshop on Prospects of Particle Physics: "Neutrino Physics and Astrophysics" Jan 26 - Ferb 2, 2014, Valday, Russia.

I. INTRODUCTION

The Baksan Neutrino Observatory INR RAS has several unique low background laboratories. One of them is deep underground low-background laboratory (DULB-4900) [1]. The laboratory is located at a distance of 3700 m from the main entry to the Baksan Neutrino Observatory tunnel, in the hall of $\sim 6 \times 6 \times 40$ m^3. Thickness of the mountain rock over DULB corresponds to 4900 m.w.e., thereby decreasing cosmic ray flux by $\sim 10^7$ times. These allow one to carry out of experiments for searching of rare process and decay. An experiment for searching of $2K$-capture of ^{124}Xe is an example.

Theoretical calculations for $2K(2\nu)$ within the frames of different models (QRPA, MCM) predict the following half-life periods with respect to $2K$-capture of ^{124}Xe: 1.08×10^{22} y [2], 3.9×10^{23} y [3]. The values were obtained by taking the fraction of $2K(2\nu)$ capture events in ^{124}Xe with respect to the total number of $2e(2\nu)$ capture events to be 73% [4].

II. BASIC ASSUMPTIONS

When two electrons are captured from the K-shell in ^{124}Xe, a daughter atom of ^{124}Te** is formed with two vacancies in the K-shell. The technique to search for this reaction is based on the assumption that the energies of characteristic photons and the probability that they will be emitted when a double vacancy is filled are the same as the sum of respective values when two single vacancies of the K-shell in two singly ionized Te* atoms are filled. In such a case, the total measured energy is $2K_{ab} = 63.62$ keV, where K_{ab} is the binding energy of a K electron in a Te atom (27.47 keV). The fluorescence yield upon filling of a single vacancy in the K-shell of Te is 0.857. The energies and relative intensities of the characteristic lines in the K series are $K_{\alpha 1} = 27.47$ keV (100%), $K_{\alpha 2} = 27.20$ keV (54%), $K_{\beta 1} = 30.99$ keV (18%), and $K_{\beta 2} = 31.7$ keV (5%) [5]. There are three possible ways for de-excitation of a doubly ionized K-shell: 1) emission of Auger electrons only (e_a, e_a); 2) emission of a single characteristic quantum and an Auger electron (K, e_a); and 3) emission of two characteristic quanta and low-energy Auger electrons (K, K, e_a), with probabilities of $p_1 = 0.020$, $p_2 = 0.246$, and $p_3 = 0.734$, respectively. A characteristic quantum can travel a long enough distance in a gas medium between the points of its production and absorption. But photoelectrons produce almost pointwise charge clusters of primary ionization in the

TABLE I. Characteristics of samples 124.

Samples, L	Xenon isotopes								
	124	126	128	129	130	131	132	134	136
	Content, L								
№ 1a, 1.98	1.02	$3.81{\cdot}10^{-1}$	$5.19{\cdot}10^{-1}$	$3.1{\cdot}10^{-4}$	$1.3{\cdot}10^{-4}$	$3.76{\cdot}10^{-5}$	$1.8{\cdot}10^{-5}$	$1.8{\cdot}10^{-5}$	$1.8e{\cdot}10^{-5}$
№ 2, 1.27	1.13	$1.3{\cdot}10^{-2}$	$3.4{\cdot}10^{-2}$	$6.1{\cdot}10^{-4}$	$1.2{\cdot}10^{-4}$	$3.6{\cdot}10^{-5}$	$3.6{\cdot}10^{-5}$	$3.5{\cdot}10^{-5}$	$3.5{\cdot}10^{-5}$
№ 3, 3.56	1.60	1.93	$1.03{\cdot}10^{-3}$	$6.8{\cdot}10^{-5}$	$6.8{\cdot}10^{-5}$	$6.7{\cdot}10^{-5}$	$6.7{\cdot}10^{-5}$	$6.6{\cdot}10^{-5}$	$6.5{\cdot}10^{-5}$
№ 4, 3.11	1.50	1.4	$1.2{\cdot}10^{-1}$	$6.7{\cdot}10^{-4}$	$6.7{\cdot}10^{-4}$	$5.8{\cdot}10^{-4}$	$5.8{\cdot}10^{-4}$	$5.8{\cdot}10^{-4}$	$5.8{\cdot}10^{-4}$
№ 5, 1.32	1.11	$4.0{\cdot}10^{-3}$	$7.5{\cdot}10^{-5}$	$5.0{\cdot}10^{-5}$	$5.0{\cdot}10^{-5}$	$3.1{\cdot}10^{-4}$	$5.0{\cdot}10^{-5}$	$5.0{\cdot}10^{-5}$	$5.0{\cdot}10^{-5}$
№ 6, 1.31	1.31	$1.6{\cdot}10^{-4}$	$1.3{\cdot}10^{-5}$	$1.3{\cdot}10^{-5}$	$1.3{\cdot}10^{-5}$	$1.3{\cdot}10^{-5}$	$1.2{\cdot}10^{-5}$	$1.2{\cdot}10^{-5}$	$1.2{\cdot}10^{-5}$

gas. In case of the event with the escape of two characteristic quanta absorbed in the working gas and a single Auger electron, the energy is distributed among three pointwise charge clusters. It is these three-point (or three-cluster) events possessing a unique set of features that were the subject of the search in Ref.[6].

To register the process of $2K$-capture in ^{124}Xe a large proportional counter (LPC) with a casing of M1-grade copper has been used [7]. The LPC is inside the shielding of 18 cm thick copper, 15 cm thick lead, and 8 cm thick borated polyethylene layers. The installation is placed in one of the chambers of the underground laboratory DULB-4900, where cosmic ray flux is lowered to the level of $(3.03 \pm 0.10) \times 10^{-9}$ cm^{-2}s^{-1} [8].

The details of the counter signals formation, of the features of pulses registration by digital oscilloscope, of the data treatment and analysis are described in Refs.[7, 9].

III. MEASUREMENT RESULTS

The measurements with radioactive-pure xenon (background sample) have been done at the initial stage of the experiment for understanding background level of experimental setup and for future comprising with ^{124}Xe. The LPC pressure is equal 1.9 atm. The measurement time was 970 hours. Count rate in interval $15 \div 150$ keV was found to be ~ 16 h^{-1}.

A sample of 12 L of xenon enriched in the isotope ^{124}Xe to 63.3% (44 g) was used in the main measurements with ^{124}Xe. This sample is combined from several one, see Tab.I.



FIG. 1. The spectra of xenon before purification (*1* - ^{124}Xe+^{85}Kr sample), radioactive-pure xenon (*2* - ^{nat}Xe sample) and xenon after purification (*3* - ^{124}Xe sample), normalized to 100 h.

The LPC was filled to the maximum pressure $P_{\max} = 1.1$ atm and the first test measurement was done during 310 hours. Count rate in interval $15 \div 150$ keV was found to be ~ 60 h^{-1}. Excess background was created by ^{85}Kr decays. This isotope is a part of natural krypton (~ 50 ppm) which dissolved in xenon.

To reduce a background of krypton an attempt was made to "wash out" of ^{124}Xe sample by "clean krypton" without ^{85}Kr. This procedure is based on the assumption that xenon will settle on a bottle wall at the liquid nitrogen temperature (-190°C) (the pressure of saturated vapor 1.93×10^{-3} mmHg) but krypton will stay in gas phase because it has a relatively large value of saturated vapor pressure (~ 1.8 mmHg). It was planned preliminary to add into the xenon a few cm^3 of the krypton and than to repeat this procedure several times. The rate of xenon treating was controlled by residual background of LPC. Unfortunately this procedure did not give any positive results. One can assume that all krypton has been absorbed by xenon similar to a charcoal trap.

A helium was chosen as an another "washing" gas. The 0.5 L of a helium was added to the bottle with the xenon per one procedure. The bottle has been frozen and helium was exhausted. The background of LPC in interval $15 \div 150$ keV became ~ 12 h^{-1}, i.e., the desired result was achieved after six procedures. Easy and chip procedure for gas cleaning from remnant of krypton was developed as a result of this work. A comparison of spectra of ^{124}Xe sample before purification (spectrum *1*), of the radioactive-pure xenon (spectrum *2*)



FIG. 2. Amplitude spectra of LPC, 1.1 atm of xenon enriched in the isotope ^{124}Xe to 63.3% (44 g): *a*) the black line - all events; the gray line - one-point events; the gray bar graph - two-point events; the black bar graph - three-point events; *b*) the three-point event spectra - full; *c*) the three-point event spectra - selected by criteria ($5.0 \leq m_0 \leq 13.0$) keV and $m_1/m_2 \leq 0.7$. The pure measurement time is 1130 h. The dashed lines indicate the boundaries of the region of interest - (63.6 ± 3.7) keV.

and of ^{124}Xe after purification - (spectrum *3*) are shown on Fig.1. All spectra are normalized to 100 hours.

The LPC was filled to the maximum possible for xenon sample pressure - 1.1 atm to perform the main measurements for search of $2K$-capture of ^{124}Xe. The efficiency of registration of two K-photons with energy ~ 27 keV is equal 0.09 at this pressure. Measurements in underground low-background setup were carried out during 1130 h. The charge pulses are recorded by digital oscilloscope to the personal computer. The collected data set was treated by special program. As a result, the entire dataset was divided into groups (spectra): one-, two- and three-point events. The full spectrum, the spectra of one-, two- and three-point events are shown on Fig.2*a* by the black line, the gray line, the gray bar graph and the black bar graph. The three-point events spectrum is shown on Fig.2*b* more precisely. The "useful" effect is searching in three-point event spectrum in energy interval (63.6 ± 3.7)

TABLE II. Characteristics of samples $N^{\underline{o}}$ 7, total vol. $\sim$ 58 L, ^{124}Xe $\sim$ (4.33 $L \approx$ 23.96 g), ^{126}Xe $\sim$ (15.24 $L \approx$ 85.7 g).

Samples, L	Xenon isotopes								
	124	126	128	129	130	131	132	134	136
	Content, L								
$N^{\underline{o}}$ 7, 58	4.33	15.24	24.147	14.033	0.0511	0.0394	0.0168	0.0606	0.0543

keV. The amplitudes of partial pulses for each event are arranged in the increased order $[(A_1, A_2, A_3) \rightarrow (m_0 \leq m_1 \leq m_2)]$, to simplify the selection of events with a given set of features. Selected useful events must satisfy the selection criteria $(5.0 \leq m_0 \leq 13.0)$ keV and $m_1/m_2 \geq 0.7$. The events corresponding to these ratios were selected from the spectrum of the three-point pulses to reduce the background. They are presented in Fig.2*c*.

The area of the expected effect marked by dashed lines. There are not remained any events in this area after the event selection. Therefore, we obtain $N_{eff} = 2.44$ for 1130 h ($\mu = 0.00 \div 2.44$) or $N_{eff} = 21.37$ y^{-1} (90% C.L.) by using of the recommendation of the work [10] for the given (effect+background) value (0 events). The half-life period has been calculated using formula:

$$\lim\ T_{1/2} \geq \ln 2 \times N \times \frac{p_3 \times \varepsilon_p \times \varepsilon_3}{N_{eff}},$$

where $N = 2.59 \times 10^{23}$ is the number of ^{124}Xe atoms in the operating volume of the counter, $p_3 = 0.735$ is a portion of $2K$-captures accompanied by the emission of two-quanta; $\varepsilon_p = 0.09$ is the probability of two K-quanta absorption in the operating volume; $\varepsilon_3 = 0.422$ is the selection efficiency for three-point events due to $2K$-capture in ^{124}Xe.

The result obtained is:

$$T_{1/2}(0\nu + 2\nu, 2K) \geq 4.6 \times 10^{20}\mathtt{y}\ (90\%\ \mathtt{C.L.}).$$

It should be noted that for the "normal" working pressure 5 atm efficiency of registration of two K-quanta is equal 0.809. A sensitivity of experimental setup to the required process at this value of efficiency is equal:

$$S = 1.46 \times 10^{21}\mathtt{y}\ (90\%\ \mathtt{C.L.}).$$

A new 58 L sample $N^{\underline{o}}$ 7 of the enriched xenon was obtained by the team at the last time. The characteristics of the sample are shown in Tab.II.



FIG. 3. Amplitude spectra of LPC with xenon sample N^{o} 7 (5 atm) - spectrum *1* and radioactive-pure xenon (5 atm) - spectrum *2*.

The LPC was filled to 5 atm by new xenon sample N^{o} 7 for checking background measurement. The spectrum (*1*) of the sample N^{o} 7 background obtained at 146 h is shown on Fig.3. The spectrum (*2*) of the radioactive-pure xenon is shown for comparison also. All spectra are normalized to 100 h. It is seen from the spectra comparison that the sample N^{o} 7 has radioactive impurities. It creates a background in a wide energy range including to the region of interest. Intense peak is visible in area of 28 keV. Differences of the spectra (*1-2*) gives a count of rate ~ 48 h^{-1} in the interval $15 \div 150$ keV. It turned out that this value decreases exponentially over time with a half-life of about 30 days. The radioactive isotope ^{127}Xe was identified as the source due to the set of the attributes. The decay scheme of isotope ^{127}Xe [11] is shown in Fig.4. The isotope ^{127}Xe can be produce by secondary cosmic rays in isotopes of sample N^{o} 7 through reaction $^{126}\mathrm{Xe}(n,\gamma)^{127}\mathrm{Xe}$ and $^{128}\mathrm{Xe}(\mu,n)^{127}\mathrm{Xe}$. There are no published data on the cross sections of the reactions mentioned above. Apparently, this can be explained by the fact that in a natural xenon content of the original isotopes is extremely small - ^{126}Xe - 0.09%, ^{128}Xe - 1.91%. Sample N^{o} 7 is highly enriched in these isotopes, and it can be used to measure the previously mentioned cross sections.

The time of about 4 months is required for decrease of ^{127}Xe background to acceptable level before using of xenon sample N^{o} 7 in the experiment for searching of $2K$-capture of ^{124}Xe. A significant amount of ^{126}Xe is present in the sample N^{o} 7. This isotope can also decay by $2K$ capture. The transition energy is 897 keV. A high limit for the $2K$-capture half-



FIG. 4. The decay scheme of isotope ^{127}Xe [11].

life of ^{124}Xe could be obtained simultaneously if the sample $N^{\underline{o}}$ 7 will be used in experiment for searching of the $2K$-capture of ^{124}Xe.

IV. CONCLUSIONS

The new experiment for searching of $2K$-capture of ^{124}Xe is done. The 12 L sample with 63.3% (44 g) of ^{124}Xe was used in measurements. The limit on the half-life of ^{124}Xe with regard to $2K(2\nu)$-capture for the ground state of ^{124}Te ($0^+ \to 0^+$, g.s.) has been found: $T_{1/2} \geq 4.67 \times 10^{20}$ y (90% C.L.).

A new 52 L sample of enriched xenon will used on the next stage of the experiment. This sample has as part of ^{124}Xe (10.6 L -58.6 g) and ^{126}Xe (14.1 L - 79.3 g). A sensitivity at 1 year measurement time of the experimental setup to the required process will be equal to

$$S = 1.46 \times 10^{21}\mathtt{y}\ (90\%\ \mathtt{C.L.}).$$

Acknowledgement. The authors gratefully thank to the E.Yu. Povolotskii for the advice on the purification of noble gases.

[1] *Gavriljuk Ju.M. et al.* // Nucl. Instr. Meth. A 2013. V7.29. P.576; arXiv:1204.6424 [physics.ins-det].

[2] *Rumyantsev O.A. and Urin M.N.* // Phys. Lett. B 1998. V.443(10). P.51.

[3] *Aunola M. and J. Suhonen J.* // Nucl. Phys. A, 1996. V.602(2). P.133.

[4] *Bambynek W. et al.* // Rev. Mod. Phys., 1977. V.49. P.49.

[5] X-Ray-Data-Booklet, Center for X-ray Optics and Advanced Light Source LBNL, http://www.scribd.com/doc/; http://xdb.lbl.gov

[6] *E.Storm and H.I.Israel,* // Atom. Data Nucl. Data Tables 1970. V.7. P.565.

[7] *Gavrilyuk Yu.M. et al.*, // Instr.Exper.Techn. 2010. V53. P.57; arXiv:0911.5403 [physics.ins-det].

[8] *Gavrin V.N. et al.* // Preprint INR RAN, P-698, Moscow 1991.

[9] *Gavrilyuk Yu.M. et al.* // Phys. Rev. C 2013. V.87. P.035501.

[10] *Feldman G.J. and Cousins R.D.* // Phys.Rev. D 1998. V.57. P.3873.

[11] *Firestone R.B.* // Table of Isotopes CD-ROM, Eighth Edition: 1999 Update, Lawrence Berkeley National Laboratory, University of California.